



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Recurrent syncope in the young: do not forget the coronary arteries

Foglia, Corinna ; Knirsch, Walter ; Valsangiacomo Büchel, Emanuela R

DOI: <https://doi.org/10.1093/eurheartj/ehs322>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-154746>

Journal Article

Published Version

Originally published at:

Foglia, Corinna; Knirsch, Walter; Valsangiacomo Büchel, Emanuela R (2013). Recurrent syncope in the young: do not forget the coronary arteries. *European Heart Journal*, 34(18):1334.

DOI: <https://doi.org/10.1093/eurheartj/ehs322>

management and it remains to be seen whether the STICH trial results trigger a paradigm shift in clinical practice.

Supplementary material

Supplementary material is available at *European Heart Journal* online.

Funding

Conflict of interest: none declared.

References

The list of references is available in the online version of this paper.

CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehs322

Online publish-ahead-of-print 16 October 2012

Recurrent syncope in the young: do not forget the coronary arteries

Corinna Foglia, Walter Knirsch, and Emanuela Regina Valsangiacomo Buechel*

Division of Pediatric Cardiology, University Children's Hospital Zurich, Steinwiesstr 75, Zurich 8032, Switzerland

* Corresponding author. Tel: +41 1 266 7339, Fax: +41 1 266 7981, Email: emanuela.valsangiaco@kispi.uzh.ch

This paper was guest edited by Brahmajee Kartik Nallamothu, University of Michigan, USA

A 9-year-old patient was resuscitated after sudden cardiac arrest (SCD) during a soccer game. The boy had a history of exercise-related syncope since the age of 5, but all the previous extensive cardiac evaluations were unremarkable.

During resuscitation, ECG documented ventricular fibrillation (*Panel A*); defibrillation was successful. At hospital admission, clinical and non-invasive cardiac findings were unremarkable. Cardiac catheterization was performed primarily for invasive electrophysiological study. Coronary angiography showed a prominent right coronary artery (*Panel B*) and the left coronary ostium was missed; all the left coronary system was exclusively perfused retrogradely (*Panel C*). Diagnosis of atresia of the left coronary artery (LCA) was done. Further non-invasive perfusion imaging was performed. SPECT with adenosine stress testing demonstrated intact myocardial perfusion (*Panel D*); CT confirmed lack of continuity between the LCA and the aortic root (*Panel D*). MRI ruled out presence of scars. LCA atresia was confirmed intraoperatively, as ostium cannulation was not possible either from the aortic root or from the LCA (*Panel E*). Surgical revascularization consisted of LIMA/LAD bypass. 3 months later, the patient was well and asymptomatic, and the aorto-coronary bypass patent.

Even though generally a rare condition, congenital anomalies of the coronary arteries are a common cause of SCD in young individuals. Among all anomalies, atresia of one coronary artery is exceedingly rare, having been reported in few isolated cases. In patients with a typical history of exercise-related ischaemic symptoms, the level of suspicion should be high and invasive evaluation initiated. Timely diagnosis is crucial for preventing SCD and planning surgical revascularization.

